



THE CITY OF MARYSVILLE DISCLAIMS ANY WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS OF THIS DATA FOR ANY PARTICULAR PURPOSE, EITHER EXPRESSED OR IMPLIED. NO REPRESENTATION OR WARRANTY IS MADE CONCERNING THE ACCURACY, CURRENCY, COMPLETENESS OR QUALITY OF DATA DEPICTED. ANY USER OF THIS DATA ASSUMES ALL RESPONSIBILITY FOR USE THEREOF, AND FURTHER AGREES TO HOLD THE CITY OF MARYSVILLE HARMLESS FROM AND AGAINST ANY DAMAGE, LOSS, OR LIABILITY ARISING FROM ANY USE OF THIS DATA.



Coho Salmon Distribution
Quilceda-Allen Watershed
 Streamnet.org Dec., 2014 Download.
 Compiling Agency: Washington Dept. of Fish and Wildlife.



TIMING OF COHO FRESHWATER LIFE STAGES

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Spawning												
Incubation												
Fry												
Junveniles												

Coho salmon utilize shallow reaches and small streams for spawning. The young spend one to two years in fresh water before migration.

- Basins**
- Allen Cr.
 - Ebey Slough
 - West Fork Quilceda Cr.
 - Edgecomb Cr.
 - West Fork Quilceda Cr.
 - Quilceda Cr.
 - Sunnyside Cr.
- Coho Use**
- Use Type**
- Migration only
 - Rearing and migration
 - Spawning and rearing